

Healthy Smile Tips

- Avoid brushing surgical area, use the ultra-soft toothbrush.
- Clean between your teeth once a day.
- Eat a healthy soft diet and limit snacks.
- Visit the dentist at your appointment.

For more information about taking care of your mouth and teeth, visit the hospital's website,



www.dentalhospitalbangkok.com

DENTAL TREATMENT

Tooth Extraction: Post-Operative Instructions



DENTAL HOSPITAL
SUKHUMVIT 77 BANGKOK

A Few Simple Guidelines

Sometimes, teeth need to be removed due to decay, disease or trauma. Having a tooth "pulled" is called a tooth extraction.

When you have an extraction, it's natural that changes will occur in your mouth. The dentist may give you instructions to follow after the extraction, and it's important to talk to the dentist if you have any questions or problems. Here are some general guidelines to help promote healing, prevent complications, and make you more comfortable.

Anesthetics

Before the extraction, you will be given an anesthetic to reduce your discomfort. Your mouth will remain numb for a few hours after the extraction. While your mouth is numb, you'll want to be careful not to bite your cheek, lip or tongue. Do not eat any foods that require chewing while your mouth is numb. The numbness should go away within several hours, if it doesn't, contact the hospital.

Bleeding

The dentist may place a gauze pack on the extraction site to limit bleeding. This will also help a blood clot to form, which is necessary for normal healing. This gauze pack should be left in place for 45 to 60 minutes after you leave the hospital.

Do not chew on the pack. There may be some bleeding or oozing after the pack is removed.

If so, here's what to do:

- Fold a piece of clean gauze into a pad thick enough to bite on. Dampen the pad with clean, warm water and place it directly on the extraction site.
- Apply pressure by closing the teeth firmly over the pad. Maintain this pressure for about 60 minutes. If the pad becomes soaked with blood, replace it with a clean one.
- Do not suck on the extraction site or disturb it with your tongue.
- Do not rinse your mouth with too much force.
- Do not drink through a straw.
- A slight amount of blood may leak from the extraction site until a clot forms. However, if heavy bleeding continues, call the hospital. (Remember, though, that a little bit of blood mixed with saliva can look like a lot of bleeding.)



DENTAL HOSPITAL

7 SOI ON NUT 1/1, SUKHUMVIT 77
SUKHUMVIT ROAD, WATTHANA
BANGKOK 10110, THAILAND
Tel. 02 092 2000
dental@dhbkk.com

Do Not Disturb!

The blood clot that forms in the tooth socket is an important part of the normal healing process. You should avoid doing things that might disturb the clot. If the blood clot is disturbed and breaks down, you can get a dry socket. Dry sockets can be extremely painful. To lower your risk of a dry socket, be very careful to not do anything that can disturb the clot.

- Do not rinse your mouth vigorously, or drink through a straw for 24 hours. These activities create suction in the mouth, which could loosen the clot and delay healing.
- Avoid alcoholic beverages or mouthwash containing alcohol for 24 hours.
- If you are a smoker, talk to your dentist before the surgery on ways to quit. You should not smoke after surgery.
- Limit strenuous activity for 24 hours after the extraction. This will reduce bleeding and help the blood clot to form.
- Sometimes the blood clot does not form in the first day or two after the extraction, or it forms but breaks down. The result is called dry socket. This can be very painful and should be reported to the dentist. A dressing may be placed in the socket to protect it until the socket heals and to reduce any pain.

**PLEASE
CALL THE
HOSPITAL**

Cleaning Your Mouth

Do not clean the teeth next to the healing tooth socket for the rest of the day. You should, however, brush and floss your other teeth well and begin cleaning the teeth next to the healing tooth socket the next day. You can also brush your tongue. This will help get rid of the bad breath and unpleasant taste that are common after an extraction.

The day after the extraction, gently rinse your mouth with warm salt water (half a teaspoon salt in an 8 oz. glass of warm water) after meals to keep food particles out of the extraction site, and try not to rinse your mouth vigorously, as this may loosen the blood clot. If you have hypertension, discuss with your dentist whether you should rinse with salt water. Avoid using a mouthwash during this early healing period unless the dentist advises you to do so.

Medication

If the dentist has prescribed medicine to control pain and inflammation, or to prevent infection, use it only as directed. If the pain medication prescribed does not seem to work for you, do not take more pills or take them more often than directed.

Swelling and Pain

After a tooth is removed, you may have some discomfort and notice some swelling. This is normal. To help reduce swelling and pain, try applying an ice bag or cold, moist cloth to your face. The dentist may give you specific instructions on how long and how often to use a cold compress.

When to call the hospital

If you have any of the following issues, call the hospital immediately.

If you cannot reach the hospital, please go to a general hospital emergency room.

- Fever, nausea or vomiting
- Ongoing or severe pain, swelling, or bleeding
- Pain that gets worse with time instead of better

AFTER THE EXTRACTION, DRINK LOTS OF LIQUID AND EAT SOFT, NUTRITIOUS FOODS. AVOID HOT LIQUIDS AND ALCOHOLIC BEVERAGES.

Eating and Drinking

After the extraction, drink lots of liquids and eat soft nutritious foods. Avoid hot liquids and alcoholic beverages. Do not use a straw. Begin eating solid foods the next day or as soon as you can chew comfortably. For the first few days, try to chew food on the side opposite the extraction site. When it feels comfortable, you should resume chewing on both sides of your mouth.

Follow-Up

If you have sutures that require removal, the dentist will tell you when to return to the hospital.

